

Dear Editor,

I would like to express my gratitude to you and your reviewers for your helpful criticism.

The major changes that we made in our article in accordance with your recommendations are as follows:

Reviewers' comments and our response to the comments:

Reviewer Number	Original comments of the reviewer	Reply by the author(s)
1	The number of doses of the subjects should be indicated instead of using the term "fully". Currently several countries are already in the 2 booster dose.	Done. "The clinical characteristics of at least two doses of vaccinated hospitalized COVID-19 patients in a Tertiary Care Hospital in Turkey"
	Improve the introduction, it's too short. There are several topics that are not contextualized for the research question.	Done.
	Give context of vaccines, types of vaccines and vaccination in the world in a paragraph.	Done.
	Should the vaccination conditions be indicated, or were they universal?	In Turkey
	Improve the discussion. It is necessary to compare their main results and consider the current situation of several countries with more doses of vaccination.	Done.
	Avoid repeating data already indicated in results.	We thought it would be good to briefly summarize the study's results at the beginning of the discussion section.
	It is suggested to transform into two tables, with the last 4 in a different table and doing multivariate analyses.	We split two tables. A multivariate binary logistic regression analysis was used to determine the influence of age, gender, presence of comorbidity, variants, vaccine status, and the time between

		the last vaccine and illness onset on the probability of mortality.
--	--	---

Sincerely, Corresponding Author,

Ayşin Kılınç Toker

Kayseri City Hospital, Department of Infectious and Clinical Microbiology/ Kayseri, Turkey.

Tel: +90 352 315 77 00

e-mail: dr.aysin@gmail.com